

CHEST[®]

Official publication of the American College of Chest Physicians



Risks in Using Nonrigorous Spanish Translations of Asthma Questionnaires

Rafael Narvaez, Kathleen Moors, Rachel Miller and Mayra Ramirez

Chest 2007;131:1271-1272

DOI 10.1378/chest.07-0180

The online version of this article, along with updated information and services can be found online on the World Wide Web at:

<http://chestjournal.chestpubs.org/content/131/4/1271.full.html>

Chest is the official journal of the American College of Chest Physicians. It has been published monthly since 1935. Copyright 2007 by the American College of Chest Physicians, 3300 Dundee Road, Northbrook, IL 60062. All rights reserved. No part of this article or PDF may be reproduced or distributed without the prior written permission of the copyright holder.
(<http://chestjournal.chestpubs.org/site/misc/reprints.xhtml>)
ISSN:0012-3692



Table 1—Rating of Clinical Areas for Quality of Training During Fellowship, Importance in Current Work Position, and Priority for Improvement in Training

Clinical Areas	Column A, Clinical Training Quality, Mean Response* (Rank Order)	Column B, Importance in Current Position, Mean Response* (Rank Order)	Column C, Training Gap (Column B Minus Column A, Difference (Rank Order)	Column D, Top Five Training Priorities†, % of Respondents Ranking This Clinical Area (Rank Order)	Column E, Training Priority Score, Sum of Rank in Columns B, C, and D (Rank Order for This Cumulative Score)	Column F, Technical Training Quality, Mean Response* (Rank Order)
Arterial blood gases	4.60 (1)	4.53 (1)	− 0.07 (13)	26.7 (7)	21 (6)	4.12 (2)
Bronchoscopy	4.50 (2)	4.37 (2)	− 0.13 (14)	43.3 (5)	21 (6)	4.17 (1)
Ventilator management	4.42 (3)	4.43 (3)	+ 0.01 (9)	57.7 (3)	15 (2)	4.08 (3)
Spirometry	4.21 (4)	4.29 (4)	+ 0.08 (7)	39.1 (6)	17 (3)	3.63 (4)
Bronchodilator response	4.17 (5)	4.03 (5)	− 0.14 (15)	9.4 (15)	35 (14)	3.54 (5)
Diffusing capacity	3.91 (6)	3.91 (6)	0.00 (10)	16.4 (12)	22 (8)	3.29 (6)
Lung volumes, gas dilution	3.86 (7)	3.64 (9)	− 0.22 (16)	8.8 (16)	41 (15)	3.26 (7)
Lung volumes, body box	3.70 (8)	3.65 (8)	− 0.05 (11)	21.0 (10)	29 (11)	3.15 (8)
Lung compliance	3.59 (9)	3.62 (10)	+ 0.03 (8)	18.8 (11)	29 (11)	3.15 (8)
Airway challenge testing	3.41 (10)	3.55 (12)	+ 0.14 (6)	21.4 (9)	27 (10)	2.99 (11)
Respiratory muscle strength	3.39 (11)	3.58 (11)	+ 0.19 (5)	23.4 (8)	24 (9)	3.02 (10)
Airways resistance	3.38 (12)	3.32 (16)	− 0.06 (12)	14.8 (14)	42 (16)	2.95 (13)
Sleep-disordered breathing	3.09 (13)	3.73 (7)	+ 0.64 (2)	52.5 (4)	13 (1)	2.90 (14)
Cardiopulmonary exercise testing	3.17 (14)	3.54 (13)	+ 0.37 (4)	61.9 (1)	18 (5)	2.99 (11)
Six-minute walk/step/ shuttle testing	2.92 (15)	3.39 (15)	+ 0.47 (3)	16.0 (13)	31 (13)	2.70 (15)
Pulmonary rehabilitation	2.74 (16)	3.52 (14)	+0.78 (1)	58.3 (2)	17 (4)	2.57 (16)

*1 = inadequate, 2 = fair, 3 = adequate, 4 = good, and 5 = excellent.

†Subjects were asked, “If you could improve the quality of training for current trainees in five of the areas, which would you choose? Please choose only five, and rank 1 through 5 with 1 being your first choice.” This column indicates the percentage of respondents including the clinical area among their top five.

American hybrid. *Thorax* 1999; 54:286–287

3 Phillips B, Collop N, Goldberg R. Sleep medicine practices, training, and attitudes: a wake-up call for pulmonologists. *Chest* 2000; 117:1603–1607

4 Angus DC, Kelley MA, Schmitz RJ, et al. Current and projected workforce requirements for care of the critically ill and patients with pulmonary disease: can we meet the requirements of an aging population. *JAMA* 2000; 284:2762–2770

5 Kvale PA, Wagner PD, Epstein LJ. Pulmonary physicians in the practice of sleep medicine. *Chest* 2005; 128:3788–3790

Risks in Using Nonrigorous Spanish Translations of Asthma Questionnaires

To the Editor:

Taking into account the substantial growth occurring among Latino populations in the United States, and the fact that some Latino populations are especially vulnerable for asthma, correctly identifying asthma cases and symptoms among persons in these populations is critical. We evaluated potential problems inherent

to the English-Spanish translation of asthma questionnaires, in particular the word “wheeze,” which is key to the performance of most of these instruments.¹

Two groups of participants were recruited (60 bilingual [Spanish-English] and 40 Spanish monolingual; all were New York residents who were parents of children with asthma-related respiratory problems). Institutional review board clearance was obtained. Bilingual participants were asked to (a) translate the word “wheeze” to Spanish, and (b) to retrotranslate a nonrigorous Spanish translation of the word “wheeze” (“ronquido”). This translation was furnished by an accredited translator who was experienced in medical translation. The translator was familiar with the vernacular of the population in the area where such nonrigorous translation was tested, which was a predominantly Dominican neighborhood in New York City that is affected by asthma rates substantially greater than those reported nationally.² This nonrigorous translation was also checked for accuracy by a second accredited translator. Monolingual participants were asked to describe asthma symptoms in Spanish. As expected, due to the local demographics of the recruitment area, the majority of our participants (84 of 100 participants) were Dominican.

The word “wheeze” presented problems for bilingual participants whose primary language was English; 9 of 39 of these respondents (23%) could not translate “wheeze” into Spanish.

The 30 participants who translated it provided 12 different translations for this word (eg, "tight chest," "whistle in the chest," "congested breathing," "hoarseness," "asphyxiation," "asthma," "snoring," and "suffocation"). Only two of these respondents (7%), agreeing with the nonrigorous translation, translated "wheeze" as "ronquido." In general, the nonrigorous translation fared poorly among most bilingual participants; 10 of 39 primary English speakers (26%) could not translate it back into English. The 29 participants who translated it back into English provided 12 different translations for the word "ronquido" (eg, "snore," "phlegm," "raspy," "hoarse," "cough," "whistling," "congestion," and "wheeze"). Only 6 of 39 of these respondents (15%), agreeing with the nonrigorous translation, translated "ronquido" as "wheeze."

The nonrigorous translation fared worse among respondents whose primary language was Spanish, as none of the 21 respondents agreed with it. The word "wheeze" also presented problems for this group. Six respondents could not translate the word into Spanish. The remaining 15 respondents provided six different translations. Responses from monolingual respondents describing asthma symptoms also were wide-ranging. After conducting this survey, a third certified translator, who was blinded to our results and was also based in the recruitment area, was consulted. She suggested that the more technical translation of "wheeze" is not "ronquido," but "sibilancia." However, none of the 60 bilingual respondents we interviewed translated "wheeze" as "sibilancia"; and none of the 40 monolingual respondents used this term to describe asthma symptoms.

Combined, these results suggest that it is quite difficult to define a single best word, in vernacular Spanish, to translate "wheeze." These difficulties inherent in cross-linguistic (*ie*, Spanish-English) measurements of asthma may have implications for asthma research that relies on responses to wheeze-related

questions. Our observations suggest that nonrigorous Spanish translations (defined as relying on one accredited translator, and neither tested by retrotranslations nor piloted among targeted cohorts) may lead to the misinterpretation of asthma questionnaires and diagnoses. Rigorous translation may be a necessary standard in the field.

Rafael Narvaez, PhD-ABD

Kathleen Moors

Rachel Miller, MD

Columbia University

New York, NY

Mayra Ramirez, RPAC, MA

Pediatrics 2000

New York, NY

The authors have reported to the ACCP that no significant conflicts of interest exist with any companies/organizations whose products or services may be discussed in this article.

Reproduction of this article is prohibited without written permission from the American College of Chest Physicians (www.chestjournal.org/misc/reprints.shtml).

Correspondence to: Rafael Narvaez, Columbia University, Medicine College of Physicians and Surgeons, PH8E, 630 W 168th St, New York, NY 10032; e-mail: rjn2101@columbia.edu

DOI: 10.1378/chest.07-0180

REFERENCES

- 1 Torén K, Brisman J, Järholm B. Asthma and asthma-like symptoms in adults assessed by questionnaires. *Chest* 1993; 104:600–608
- 2 Crain EF, Weiss KB, Bijur PE, et al. An estimate of the prevalence of asthma and wheezing among inner-city children. *Pediatrics* 1994; 94:356–362

Risks in Using Nonrigorous Spanish Translations of Asthma Questionnaires

Rafael Narvaez, Kathleen Moors, Rachel Miller and Mayra Ramirez

Chest 2007;131; 1271-1272

DOI 10.1378/chest.07-0180

This information is current as of February 1, 2011

Updated Information & Services

Updated Information and services can be found at:

<http://chestjournal.chestpubs.org/content/131/4/1271.full.html>

References

This article cites 2 articles, 1 of which can be accessed free at:

<http://chestjournal.chestpubs.org/content/131/4/1271.full.html#ref-list-1>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://www.chestpubs.org/site/misc/reprints.xhtml>

Reprints

Information about ordering reprints can be found online:

<http://www.chestpubs.org/site/misc/reprints.xhtml>

Citation Alerts

Receive free e-mail alerts when new articles cite this article. To sign up, select the "Services" link to the right of the online article.

Images in PowerPoint format

Figures that appear in *CHEST* articles can be downloaded for teaching purposes in PowerPoint slide format. See any online figure for directions.

